# This Page Is Inserted by IFW Operations and is not a part of the Official Record

# **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

# IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.





Patent Office Canberra

I, JULIE BILLINGSLEY, TEAM LEADER EXAMINATION SUPPORT AND SALES hereby certify that annexed is a true copy of the Provisional specification in connection with Application No. 2002950182 for a patent by PETER JAMES JENKINS as filed on 12 July 2002.

WITNESS my hand this Second day of February 2004

JULIE BILLINGSLEY

TEAM LEADER EXAMINATION

**SUPPORT AND SALES** 

## **Compounds For Medicinal Purposes**

#### Abstract:

Application of compounds of formula (1) (Gibberellins) and their derivatives for the preparation of a pharmaceutical composition or medicaments for the treatment of diabetes and related conditions. The method results the normalization of serum glucose level and other physiological conditions.

#### Field of the invention:

The present invention relates to the application of a group of compounds known as Gibberellins and their derivatives for the preparation of a pharmaceutical composition for the treatment of diabetes and related conditions, as well as a method for treating these and other conditions by administering Gibberellins on a pharmaceutically acceptable salts or esters including glycoside esters, active esters or lactones. Moreover, this invention relates to the manufacturing and the use of a medicament for treating diabetes and related conditions thereof. Furthermore, the application of Gibberellins and their derivatives especially by oral, injection, transdermal patches, or by inhalation administration can be used as a substitute for insulin and/or its fragment derivatives and/or IGF (Insulin like Growth Factor) treatment or as a choice of combination therapy with insulin, its fragment derivatives, IGF, growth factors or other pharmaceutically compatible anti-diabetic agents for the treatment of diabetes and related conditions.

### Background of the invention:

This invention relates to a novel application of Gibberellins in veterinary and human medicines. In particular the invention concerns Gibberellins' pharmaceutical formulations and their use for the treatment of diabetes including type 1 and type 2 diabetes and their related conditions.

Gibberellins are a series of naturally occurring compounds, which are known as plant growth regulators with wide application in the plant kingdom [1]. They have also been isolated from metabolites of some microorganisms, such as Gibberella fujikuroi [2]. Gibberellins, especially Gibberellic Acid (Gibberellin A<sub>3</sub>), and its mixture with Gibberellin A<sub>4</sub> and/or A<sub>7</sub> which are commercially available, have been extensively applied in agriculture to increase the growth of some fruits (strawberries and grapes) and

vegetables (tomatoes, cabbages and cauliflowers), also as food additive in the malting of barley [3].

[1]. J. MacMillian, et al. "Isolation and Structure of Gibberellin From Higher Plants". Adv. Chem. Ser 28, 18~24, (1961).

[2].

- (a) P.J. Curtis et al. Chem. & Ind. (London) 1066, (1954).
- (b) B.E. Cross, J. Chem. Soc. 4670, (1954).
- (c) P.W. Brian et al, U.S. 2,842,051.
- (d) C.T. Calam et al, U.S. 2,950,288.
- (e) A.J. Birch et al, U.S. 2,977,285.

[3].

- (a) M. Devlin, Plant Physiology, New York, Reinhold, (1966).
- (b) P.W. Brian et al, Plant Physiol, 5,669 (1955).
- (c) A.K Mehta et al, J. Hostic Sci 4, 167 (1975).
- (d) R.J. Weavor, Adv. Chem. Ser 28, 89 (1961).
- (e) F.G. Gustafson, Plant Physical 35, 521 (1960).
- (f) Fed. Reg. 25, 2162 (1960).

Gibberellin  $A_3$  and its mixture of Gibberellin  $A_4$  and/or  $A_7$  can be obtained by fermentation of microorganisms such as Gibberella fujikuroi. The crude compounds can be isolated and purified by solvent extraction and recrystallization to afford a high purity crystalline product. The other derviatives of Gibberellin can be obtained by either semi-synthetic route from Gibberellin  $A_3$  or total synthesis which have been well documented [4].

[4].

- (a) The Merck Index, 12, 4426, literatures cited herein.
- (b) Furber M., et al., "New Synthesis Pathways From Gibberellins to Autheridiogens Isolated From the Fern Genus Anemia", J. of Org. Chem. vol 55, No. 15, 4860~4870 (1990).
- (c) Mander L. N., et al., "C-18 hydroxylation of Gibberellins", J. C. S., Perkin Trans. 1 (17), 2893~2894 (2000).
- (d) Pour M. et al., "Synthesis of 3,12-Dihydroxy-9,15-Cyclo Gibberellins", Tetrahedron 54(45), 13833~13850 (1998).
- (e) Liu J. P. et al., "A General Protocol For the Hydroxylation of C-14 in Gibberellins Synthesis of 14-Beta-hydroxy-Gibberellin A<sub>1</sub> Methyl Ester", Tetrahedron 54(38), 11637~11650 (1998).
- (f) Pour M, et al., "Synthetic and Structural Studies on Novel Gibberellins", Pure and Applied Chemistry 70(2), 351~354 (1998); "Synthesis of 12-Hydroxy-9,15-Cyclo-Gibberellins", Tetrahedron Letters, 39(14), 1991~1994 (1998); Australian J. of Chemistry 50(4), 289~299 (1997).

- (g) King G. R. et al., "A New and Efficient Strategy for the Total Synthesis of Polycyclic Diterpenoids The Preparation of Gibberellins (+/-)-GA<sub>103</sub> and (+/-)-GA<sub>73</sub>", J. Am. Chem. Soc. 119(16), 3828~3829 (1997).
- (h) Mander L. N., "Synthsis of 12-Hydroxy-C-20-Giebberellin from Gibberellin A<sub>3</sub>", Tetrahedron 53(6), 2137~2162 (1997) and literatures cited herein.

Furthermore, the extraction and isolation of different Gibberellins from different plants, shoots, fruits and seeds have also been widely published [5].

- [5].
- (a) Pearce D.W., et al., Phytochemistry, 59(6), 679~687 (2002).
- (b) Chang S. T., et al., Physiologia Plantarum, 112(3), 429~432 (2001).
- (c) Nakayama M. et al., Phytochemistry, 57(5), 749~758 (2001); 48(4), 587~593 (1998).
- (d) Blake P. S., et al., Phytochemistry, 55(8), 887~890 (2000); 53(4), 519~528 (2000).
- (e) Koshioka M., et al., J. of the Japanese Society for Horticultural Science, 68(6), 1158~1160 (1999); 67(6), 866~871 (1998).
- (f) Mander L. N. et al., Phytochemistry, 49(8), 2195~2206 (1998); 49(6), 1509~1515 (1998).
- (g) Wynne G. et al., Phytochemistry, 49(7), 1837~1840 (1998).

Gibberellins have previously been used for anti-inflammation, treatment of prostatitis and psoriasis, treatment of tumor, and for ulcer and wound healing [6].

- (a) U.S. 4424232 1/1984 Parkinson
- (b) French 2597339 10/1987
- (c) U.S. 5487899 1/1996 Davis
- (d) U.S. 5580857 12/1996 Oden
- (e) AUS. 695054 11/1998 Wu
- (f) U.S. 6121317 9/2000 Wu

We have now found application of Gibberellin or its derivatives for the treatment of diabetes including type 1 and type 2 diabetes and their related conditions.

#### Disclosure of the invention:

It has now been found that Gibberellins possess mammalian growth factor (such as IGF, EGF) like properties in our laboratory.

The experimental results (examples 3 and 4) suggested that Gibberellins, which are generated by plants and microbes, act as broad spectrum binders binding to a range of growth factor receptors. They differ from the growth factors found in animals, each of which has a high affinity for a specific receptor. This is the result of evolution. The

biological systems of plants and microbes produce biological substances acting on a broader (less specific) base than that of the more complex life forms such as animals.

Since Gibberellins are smaller molecules than growth factors, the binding of Gibberellins on the growth factor receptors is probably weaker. In the presence of low level of growth factors, Gibberellins bind to growth factor receptors to stimulate cell growth and other functions. Under this condition, Gibberellins perform the functions of the growth factors. In the presence of normal level of growth factors, the growth factors bind to their receptors more readily due to their higher affinity for those receptor sites. The physical bulkiness of these growth factors leave no room or very little room at the receptor sites for which Gibberellins can bind. This results in Gibberellins being ineffective when growth factors are present in sufficient quantities. This mechanism provides a very good profile for Gibberellins acting as a substitute for growth factors including IGF since the presence of excess Gibberellins will not interfere with the normal functions of these growth factors.

It is known that insulin, IGF and EGF receptors are all in the same family and their structures are expected to be have 90% similarity (Ward C.M. et al., Nature, 394 (6691), 395~399 (1998); Molecular pathology, 54(3), 125~132 (2001)). Therefore, it may be logically expected that Gibberellins could play a role not only as a substitute for growth factors but also as a substitute for insulin.

Diabetes mellitus is a chronic disorder manifested by hyperglycemia and altered lipid and protein metabolism. According to the American Diabetes Association, more than 13 million people in the U.S. suffer from diabetes, and each year some 650,000 new cases are identified. The introduction of insulin and of sulfonyl ureas represented important landmarks in the treatment of diabetes mellitus. Insulin like growth factor - 1 (IGF-1), a molecule with structure homology to insulin, has its own specific receptor, the type-1 IGF receptor, through which it elicits a variety of metabolic effects that are similar to insulin. The discovery of the active region of human growth factor responsible for the insulin like actions of the molecule has led to the development of its fragment as new anti-diabetic peptide agent. It is also known that growth factors are polypeptides that regulate the replication, differentiation and metabolic homeostasis cells. They increase the growth and/or survival of neurons. IGF-2 is known to increase the rate of nerve regeneration, in pre-clinical testing for the treatment of various neurological disorders including diabetic neuropathies. Furthermore, elevated intracellular concentrations of c-AMP potentiate glucose-dependent insulin secretion from pancreatic β-cells. It is known that Gibberellins increase the activity of adenylate and gryanylate cyclase. The intracellular concentrations of c-AMP and c-GMP may therefore be increased by the administration of Gibberellins as the consequence to potentiate glucose-dependent insulin secretions from pancreatic β-cells. From the review described above, it may be logically expected to apply Gibberellins for the treatment of diabetes.

The animal experiment results (examples 5 & 6) showed diabetic rats treated with 5mg/kg of Gibberellin A<sub>3</sub> or a mixture of A<sub>3</sub> and A<sub>4</sub> or A<sub>7</sub> returned their serum glucose level to the normal, as well as their body weights. It indicated that Gibberellins may be effective in the treatment of diabetes.

In combination with the fact that the toxicity to mammals of Gibberellin A<sub>3</sub> is extremely low. The acute oral LD<sub>50</sub> for rats and mice is reported to be 6.3g/kg [7a] and >15g/kg [7b] respectively. In 90-day feeding trials, the no effect level for rats and dogs was >1g/kg/day [7b]. It is non-irritating to skin and eyes [7b]. No indication has been found of carcinogenicity [7c]. Classifications: WHO Toxicity Class Table 5 (least hazardous class product, unlikely to present acute hazard in normal use); EPA Toxcity class III (second least hazadous classification).

- (a) NTP Chemical Repository, http://ntpserver.niehs.nih.gov/htdocs/CHEM\_H&S/NTP\_CHEM7/Radian77-06-5.html
- (b) The Agrochemicals Handbook, Royal Society of Chemistry, August 1991.
- (c) Gold L. S., Slone T. H., Ames B. N. (2001), Pesticide Residues in Food and Cancer Risk: A Critical Analysis, Publications from the Carcinogenic Potency Project, in Handbook of Pesticide Toxicity, Second Edition, (R. Krieger, ed.), Academic Press.

Thus this invention provides an aspect use of compounds of formula (1) (Gibberellins) and their derivatives for the treatment of diabetes and related conditions. Furthermore, this invention would be extended as described in the "Field of the invention".

wherein

 $R^1$  is H or a group  $-O-R^{20}$ , where  $R^{20}$  is H, or together with  $R^2$  or  $R^{10}$  forms a bond ( $C_1-C_2$  or  $C_1-C_{10}$  double bond, respectively);

 $R^2$  is H or a group -O- $R^{21}$ , where  $R^{21}$  is H, a glycosylic ether group (glycoside ether) or together with  $R^4$  forms a bond (lactone) or together with  $R^1$  or  $R^3$  forms a bond (C<sub>1</sub>-C<sub>2</sub> or C<sub>2</sub>-C<sub>3</sub> double bond, respectively);

 $R^3$  is H, =O, or -O- $R^{22}$ , where  $R^{22}$  is H or a glycosylic ether group (glycoside ether), or together with  $R^2$  forms a bond ( $C_2$ - $C_3$  double bond);

 $R^4$  is OH, or  $-OR^{23}$ , where  $R^{23}$  is unsubstituted or substituted  $C_{1\sim20}$  alkyl, allyl, aryl, arylalkyl, amidine,  $-NR^{24}R^{25}$  or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulfur;  $R^{24}$  and  $R^{25}$  may or may not be the same, are hydrogen, or  $C_{1\sim20}$  alkyl, allyl, aryl, arylalkyl or an unsaturated or saturated ring containing one or more hetero-atoms slected from the group consisting of nitrogen, oxygen and sulphur;  $R^4$  together with  $R^{21}$  or  $R^{28}$  forms a bond (lactone);

 $R^5$  is H or a glycosylic ester (glycoside ester) group, or unsubstituted or substituted (e.g. haloginated)  $C_{1\sim20}$  alkyl esters, allyl esters, aryl esters, arylalkyl esters, active esters (such as phenacyl ester, pivaloyl ester);

R<sup>6</sup> is H or OH or together with R<sup>7</sup> forms a bond (C<sub>11</sub>-C<sub>12</sub> double bond);

 $^{\circ}$  R<sup>7</sup> is H, =O, or  $-OR^{26}$ , where R<sup>26</sup> is H or a glycosylic ether group (glycoside ether) or together with R<sup>6</sup> forms a bond (C<sub>11</sub>-C<sub>12</sub> double bond);

 $R^8$  is H, hydroxyl, mercaptan, or halogen (e.g. F, Cl), amino, azido,  $NR^{24}R^{25}$ , unsubstituted or substituted (e.g. halogenated)  $C_{1\sim20}$  alkyl, allyl, aryl, or arylalkyl, or  $-CR^{27}$ , where  $R^{27}$  is a glycosylic ether group (glycoside ether);

R<sup>9</sup> is H or OH, or together with R<sup>15</sup> forms a bond (C<sub>9</sub>-C<sub>15</sub> bond);

 $R^{10}$  is H, CH<sub>3</sub>, CHO, COOH, or a glycosylic ester (glycoside ester) of said COOH, CH<sub>2</sub>O-R<sup>28</sup> or -OR<sup>28</sup>, where R<sup>28</sup> is H or together with R<sup>4</sup> forms a bond (lactone) or together with R<sup>1</sup> forms a bond (C<sub>1</sub>-C<sub>10</sub> double bond);

R<sup>11</sup> is OH or absent,

R<sup>12</sup> is CH<sub>3</sub>, CH<sub>2</sub>OH, COOH or a glycosylic ester (glycoside ester) of said COOH;

 $R^{13}$  is methylene, or a divalent hetero-atom, or  $NR^{29}$ , where  $R^{29}$  is  $NR^{30}$  or  $OR^{30}$  where  $R^{30}$  is H, or  $C_{1\sim 20}$  alkyl, aryl, alkylaryl; when  $R^{11}$  is absent, a double bond is present between  $C_{16}$  and  $R^{13}$ ;

R<sup>14</sup> is H or OH;

R<sup>15</sup> is H, or together with R<sup>9</sup> forms a bond (C<sub>9</sub>-C<sub>15</sub> bond);

And pharmaceutically acceptable derivatives, lactones, esters and salts including alkali metal salts (e.g.  $Na^+$ ,  $K^+$ ), alkaline earth metal salts (e.g.  $Ca^{2^+}$ ,  $Mg^{2^+}$ ), metal salts (e.g.  $Zn^{2^+}$ ,  $Al^{3^+}$ ), and salts of ammonium, organic bases (such as  $NR^{16}R^{17}R^{18}R^{19}$  where  $R^{16}$ ,  $R^{17}$ ,  $R^{18}$ ,  $R^{19}$ , which may be the same or not the same, are hydrogen,  $C_{1-20}$  alkyl, alkanol, arryl) thereof.

The dotted line together with the solid line indicate that a double bond may be situated between two of the three carbon atoms connected by the dotted and solid lines; with the proviso that a double bond is not present if R<sup>11</sup> is an OH group.

Since Formula (1) complies with normal valence rules, this leads to the further provisos as follows:

 $R^1$  and  $R^2$  cannot form a bond if  $R^{10}$  and  $R^1$  and/or  $R^2$  and  $R^3$  form a bond;  $R^{10}$  and  $R^1$  cannot form a bond if  $R^{10}$  and  $R^{23}$  form a bond;  $R^2$  and  $R^1$  or  $R^2$  and  $R^3$  cannot form a bond if  $R^4$  and  $R^{21}$  form a bond.

In the case of Gibberellin  $A_3$ ,  $R^1$  together with  $R^2$  forms a bond ( $C_1$ - $C_2$  double bond);  $R^3$  is  $\beta$ -OH,  $R^4$  together with  $R^{28}$  forms a bond (lactone);  $R^5$ ,  $R^6$ ,  $R^7$ , and  $R^9$  are hydrogens,  $R^8$  is OH,  $R^{11}$  is absent;  $R^{12}$  is methyl;  $R^{13}$  is methylene, a double bond is present between  $C_{16}$  and  $R^{13}$ ;  $R^{14}$  and  $R^{15}$  are hydrogens.

#### **MD-960 - TRIAL 3**

#### **METHODS**

Male Wistar rats (290-330g) were weighed and lightly anaesthetised (4% halothane, 2:1 O<sub>2</sub>/N<sub>2</sub>0) so that blood glucose levels could be measured via a tail vein sample, using a Precision Q.I.D. glucometer. Diabetes was then induced by a single tail vein injection of streptozotocin (STZ, 60 mg/kg), which was dissolved immediately prior to use in citrate buffer (50mM citric acid and 50mM trisodium citrate; pH 4.5). An equivalent volume of citrate buffer was injected into age-matched control rats.

Rats were housed in groups of two for the next two weeks. Animal house temperature was maintained at 20°C (±2°C) with a 12 hour light/dark cycle, and rats were allowed free access to food and water.

Ethical approval for all experiments was obtained from the Pharmacology Animal Ethics Committee.

#### DRUG ADMINISTRATION AND DAILY MONITORING PROTOCOL

Forty eight hours after the administration of STZ (60mg/kg), a blood glucose sample was taken and animals with blood glucose levels ≥16mM were considered to be diabetic. Rats were then randomly divided into one of the following three groups:

- (1) insulin-treated (4U; s.c.) diabetic rats receiving insulin daily for the entire trial,
- (2) sub-maximal insulin- treated (2U; s.c.) diabetic rats receiving a daily dose of insulin for the entire trial. These animals also received an oral dose of MD-960 (5mg/kg) three times daily from days 42-46, and
- (3) sub-maximal insulin- (2U; s.c.) and MD-960-treated (5mg/kg) diabetic rats receiving a daily dose of insulin and MD-960. MD-960 was initially given s.c., but was administered i.p. from days 26-46. This group was given an additional dose of MD-960 (5mg/kg; i.p.) (ie. once in the morning and again in the afternoon) from days 38-46.

The slow-acting, Lente Monotard insulin was used and MD-960 was made up as required in distilled water immediately prior to use.

Blood glucose readings were obtained two hours after the administration of drug(s), every three days. From days 36-46 blood glucose readings were obtained five hours after the administration of insulin and/or MD-960.

Rats were sacrificed by a blow to the head and decapitation on day 46 of the trial in accordance with the ethics obtained for this study.

¥ a	
9	
4	•

27/05/02 319 339		. 2252	27/05/02 89.60674 101.497 108.3077	94,4444 92,75362 98,54726 92,0765		27/05/02			17.0 329.75 329.5 328.5		17/0 98.5747 92.92157 98.92394
341	ikan ka	2222	26/05/02 95.7852 102.0952 112.0231 105.2469	92.75362 100.2882 91.80328	105.6428 108.5174 102.2535 134	26/05/02			16/05/02 3.391165 5.18009 9.064563	26/5/02 25 2.216745 75 1.756595 1.756395	16/05/02 100.7564 1.389351 04.78988 0.929034 90.98837 0.48301
25/05/02 346 338	erer re	8858	25/05/02 97,75281 101,497 108,9231 106,4815	94.4444 93.04348 100.2882 92.0765	105.3292 107.2555 104.507 102.4815	25/05/01 4.5 13.5 11.5	25.25 25.25 25.25	15 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	222	28/ 10.925 15.575 7.7	167 100.7594 94.78988 99.99337
24/05/02 343 334	1226 E	8888	24/05/02 98.34831 100 103,6923 (04,9383	93.61111 93.33333 99.42363 90.98381	104,3887 106,9401 101,4085 103,0769	24'05/02			6/05/02 3.559026 4.743416 5 6.559381		15/05/02 22 1.840194 29 0.615989 83 0.731618
23/05/02 327	# # # # # # # # # # # # # # # # # # #	2823	23/05/02 95.78652 97.90419 104 102.1605	92.5 93.04348 96.2536 92.0765	103.7618 105.3628 101.9718 100.9231	23/05/02			328	22/06/02 15 0.96296 25 2.574353 15 2.574353	15( 98,07922 92,81229 98,03583
22/05/02 343 328	22 22 22	22 22 22	12/05/02 96.34831 97.60479 101.8462 101.5432	93.05556 91.88408 99.42363 92.34973	102,5078 105,0473 99,43862 100	22/05/02 3.6 5.8 8.3 5.6	14.5 6.6 6.2	4.4.00 80.00 4.00 80.00	1478/02 5 1.848423 15 4.871259 5 7.190747	22/C 5.625 10.625 5.825	14/05/02 132 1.80572 526 1.032418 385 1.197403
21/05/02 343 323	28 25 88 88 88	2222	21/05/02 96.34831 96.70659 101.8462 101.2346	92.77776 90.72464 97.98271 92.89817	101,5674 103,7655 98,30986 99,38462	21/05/02		٠	333. 330.2 310.		14/ 98.74432 93.17526 97.16385
20/05/02 318 300	38 88 3	2482	20/05/02 69.32584 69.82036 101.5385 102.7778	93.33333 69.56522 98.84728 93.96907	99,68652 104,1009 99,71831 100	20/05/02			13/05/02 5 5.693594 4.813176 15 8.673475	19/05/02 8 2.104757 25 2.009301 5 2.009301	13/05/02 857 0.535922 871 0.463087 127 1.318098
19/05/02 350 335	35 35 35	2222	19/05/02 98.31461 100.2994 102.4615 108.642	95 91.88406 100 94.80874	101.2539 104.4164 101.9718 103.0769	19/05/02 5.9 15.3 13.1 8.9	24 20.5 14.8 21.8	9.3 10.5 10 23.2	326.	19/1 10.8 20.225 15.5	97.56 92.52 96.45
18/05/02 336 330	8888 88	2552	18/05/02 94.30202 98.3044 99.38452	91.66567 90.43478 96.2536 92.89817	97,80564 100,6309 96,30988 99,38462	18/05/02	•		12/6/02 2.614065 5 5.123475 15 9.003472		12/5/02 116 1.601084 208 0.602605 163 1.172257
1778502 338 732	32 225 32 225	2442	17/05/02 94.94362 100.8982 100 98.45679	91.66667 89.27536 95.38905 95.35519	100.9404 100.3155 96.90141 97.53846	17/05:02			12 E	16/05/02 25 0.507239 375 2.719796 75 2.719796	25.03.24 26.03.19 27.03.19
16/05/02 344 341		32.5	16/05/02 98.62921 102.0958 102.7892 101.5433	93.33333 97.40634 94.80874	99.37304 101.2818 100.2817 99.07692	16/05/02 2.6 3.4 4.8 4.5	5.5 12.5 18.2 15.3	9.5 8 8.3	11/5/02 75 2.015564 25 3.902456 1 6.78233	167 3.825 12.875 6.575	11/6/02 101 2.210565 115 0.58862 363 0.975651
15/05/02 332 336	18 asas	32 33 2	15/05/02 93,25643 100,5988 99,07692 99,38272	91,94444 91,88406 94,5245 92,89617	98.74508 99.35909 96.02517 96	16/05/02			328		98.051 92.04(
1405/02	* * * * * * * * * * * * * * * * * * *	8 2 5 5	14/05/02 B4.302/02 100.8982 102.1538	90.27776 93.3333 95.10086 93.96907	96,8652 100,6309 85,77465 95,38462	14/05/02			10/6/02 4:554747 75 5:406401 5 8:62651	13/05/02 75 0.546771 375 4.238784 225 4.238784	10/6/02 74 2.144794 (8) 0.569379 (0) 0.473236
13/05/02 342 328		8 9 7 6	13/05/02 96.06742 98.20359 97.53646 98.45679	91.86667 91.88406 93.65994 92.89617	98.55172 98.68454 98.33603 93.23077	13/05/02 6.3 6.1 5.2 3.9	5.5 21.5 19.9 24.6	20.3 17.7 18.7 8.2	320	5.375 17.875 10.225	98.394 93.014 94.983
12/8/02 325 325	320 320	208 311 302	12/6/02 91.29213 97.30539 98.46154 96.91358	90.55556 69.27536 92.21902 90.71038	93.41693 98.10726 95.21127 92.92308	12/5/02			9.6702 5 4.871259 5 5.893148 5 7.888547		86.21441 2.142037 93.14834 0.608483 95.07611 0.492261
11/6/02 3.78 33.2		9888	11/5/02 91.57303 99.4012 101.5385 99.69136	90.63333 91.88406 93.65894 91.80328	97.17868 97.16088 94.08451 93.53848	11/5/02			331.7 330.2 312.7	10/6/02 0.403113 5 1.101798 5 1.101798	98.21441 93.14834 95.07611
10/8/02 330 338	12 8263	337 28	10/6/02 92,69663 101,1976 97,53089	92,22222 91,88406 94,23631 93,71585	93.73041 95.26814 94.92858	10/5/02 5.9 7.3 5.5 6.7	2. r o 1. 2. t	5.0 5.2 12.4	6/8/02 1 2.397916 5.24/044 7.767453	10, 6.35 9.375 0.275	8/5/02 24 1.755618 24 0.226902 16 0.444802
9/5/02 332 345	# # # # # # # # # # # # # # # # # # #	308 309	9/5/02 93.25843 103.2934 100.9231 99.38272	92,77778 91,5942 94,23631 93,88907	94.35737 96.52997 94.64789 94.78923	9/5/02			328.		8/5/ 98,2438 82,23634 93,93816
87802 331 334	18 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	32.22.25	8/6/02 92:97753 100 100 100	91,94444 92,17391 91,93064 92,69617	93,41693 95,26814 93,52113 93,53846	8/5/02			6/5/02 4.203173 i 5.074446 i 8.831761	6.1502 0.155456 0.262996 0.262996	6/5/02 0 0
\$56 356 370	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	319 355 325	6/5/02 100 100 100 100	8888	\$ <b>8 8 8</b>	6/5/02 6 6.5 6.7 6.0	2222	6004	354.5 324.5	6.45 6.75 7	\$ 8 8 8
			PASE .								
		<u> </u>	SOECR!		3333	w		3333	Freez	S C C C C C C C C C C C C C C C C C C C	10 C C C C C C C C C C C C C C C C C C C
DY WEIGHT TREATMENT INSULIN (4U) INSULIN (4U)	INSPILIN (20) INSPILIN (20) INSPILIN (20) INSPILIN (20) INSPILIN (20) INSPILIN (40)	• INSUI • INSUI • INSUI • INSUI	CREASE ATMENT LIN (4U) LIN (4U)	INSULN (2U) INSULN (2U) INSULN (2U)	NSU +	OOD GLUCOSE TREATMENT INSULIN (4U) INSULIN (4U) INSULIN (4U)	INSTITIN (20) INSTITIN (20) INSTITIN (20) INSTITIN (20) INSTITIN (20)	USU	Y WEIGH ATMEN JUN (4U JUN (2U	- BLOOD GLUCOSE TREATMENT INSULIN (4U) INSULIN (2U) MD-860 + INSULIN	DOY WEI SATMEN ULIN (AL
TRE	DSNI DSNI DSNI DSNI	MD-860 • INSULIN MD-860 • INSULIN MD-860 • INSULIN	TYEN TRE INSU INSU INSU	NSU NSU	MD-860 + INSULIN MD-860 + INSULIN MD-860 + INSULIN MD-860 + INSULIN	BLOOD GLUCOSE TREATMENT INSULN (4U) INSULN (4U) INSULN (4U) INSULN (4U) INSULN (4U) INSULN (4U)	INSI INSI INSI INSI INSI INSI INSI INSI	NTD:SO: + 098-CJW NTD:SO: + 108-CJW NTD:SO: + 108-CJW NTD:SO: + 108-CJW	A - BODY WEIGHT TREATMENT INSULIN (4U) INSULIN (2U) MD-080 + INSULIN	A - BLOA TRE INSI IND-96	A·* BK TRI INS INS MD-98
RAW DATA - BODY WEIGHT CODE TREATMEN PINK 1 INSULIN (4) PINK 2 INSULIN (4)	PLUK 4 BLUE 1 BLUE 2 BLUE 3 BLUE 3	- ~ ~ ~	BODY WEIGHT - % INCREASE/DECREASE CODE TREATMENT PINK 1 INSULU (4U) PINK 2 INSULU (4U) PINK 3 INSULU (4U) PINK 4 INSULU (4U)	BLUE 1 BLUE 2 BLUE 3 BLUE 4	GREEN 1 GREEN 3 GREEN 3	RAW DATA - 6 CODE PINK 1 PINK 2 PINK 3	9LUE 1 8LUE 1 BLUE 1 BLUE 4	GREEN 2 GREEN 3 GREEN 4	GROUP DATA - BODY WEIGHT CODE TREATMENT PHK INSULIN (41) BLUE INSULIN (21) GREEN MD-900 + INSULI	OROUP DATA-BLOOD GLUGOSE CODE TREATMENT PINK INSULIN (4U) BLUE INSULIN (2U) GREEN MD-980+ INSULIN	GROUP DATA - % BODY WEIGHT CODE TREATMENT PINK INSULIN (4J) BLUE INSULIN (2J) GREEN MD-800 - INSULIN


						21/06/02 2.1 3.4 4.8 2.2	5.1 13.9 4.8	8. 8. 4. g.	354 340.25 347	19/06/02 5 1.488568 75 2.584046 25 2.584046	30/0 105.6557 96.0346
380 380 359 372 362	352 338 318	356 379 368 359	21/06/02 106.7416 107.485 111.7284 97.77778	98.26087 102.5937 87.15847	112,2257 118,5584 109,2858 110,4815	21706/02 3.7 0.8 5 3.9	11.3 16.6 13.8	4 6 6 6 6	29/05/02 ) 3,53534 25 4,110454 25 6,381941	19/4 6.95 13.775 4.425	28/05/02 177 1.496624 584 1.22357 262 1.462583
2006602 1 386 387 387 385	3 8 8 5	8 2 2 3 8		95.65217 99.71182 84.97288	112,8527 120,8202 110,1408 111,3846	20/06/02			35 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5		28/ 104.6477 96.57584 105.0263
367 387 387 382 363	358 357 304	356 377 385 361		97,3913 102,6818 83,06011	111.5987 118.9274 108.4507 111.0769	19/06/02 3.9 4.9 9.7 9.3	13.5 15.8 6.6 19	3.8	5/02 3.162278 6.435578 6.745369	2002 0.443471 2.830589 2.830589	28/05/02 103.7613 1.953598 95.24676 2.085565 105.5573 1.573627
8006/02 1 380 358 359 359	355 333 316	380 387 387		96.52174 102.3055 88.3388	112.8527 119.8738 109.0141 109.8462	18/06/02			28/05/02 347 3.11 337.5 6.4 347 6.7	17/6/2002 5.3 0.44 19.025 2.83 7.475 2.83	28/0 103.7613 95.24678 105.5573
7706/02 1 378 356 367 357	348 325 325	25 17 19 25 17 18 18 18 18 18 18 18 18 18 18 18 18 18	17,06,02 106,4607 106,5668 112,923; 110,1652		110.6583 117.8656 110.1408 112	17/06/02 4.4 5.6 4.8 6.4	12.9 18.2 20.9 26.5	8.1 5.8 11.2	5/02 8.637397 5.147815 9.082951		5/02 4.0716 1.522641 0.219474
374 374 351 351 361 351	350 340 321	352 374 386 380		98,55072 9 103,7464 1 87,70492 8	110,3448 1 117,8811 1 108,2958 1 110,7892				27/05/02 337.5 8.6 335 5.1	2002 0.511534 1.542387 1.542387	27/05/02 101,0874 4.( 94,53046 1.5; 105,1728 0.2;
106/02 16 358 368 348	353 352 321	2		97,97101 9 104,3228 1 87,70492 8	119,9718 1 119,5584 1 107,0423 1 109,2308 1	15/06/02 16/06/02			6.5 6.5 5.764475 6.034622	14/6/2002 3.7 0.51 10.175 1.54 9.125 1.54	3.5621 1.927073 1.331826
106/02 11 376 357 357 358	350 335 321	355 376 391		97.10145 9 103.4562 1 87.70492 8	111,2853 1 116,812 1 110,1408 1 110,7692 1	3.1 3.1 5 4 2.7	9.7 14.5 7.2 9.3	10.8 3.5 11.4 10.8	2605/02 347.5 6 335.25 5.76 346.5 6.00		26/0 104.0131 94.61405 105.1034
377 377 352 362 358	32 32 52	385 374		98,26087 9 101,7291 1 88,25137 8	109,4044 1 117,9811 108,4507 1 109,5365 1	13/06/02			3.122499 5.664215 8.784456	5/02 0.488834 2.045065 2.045085	702 2.503899 1.840356 0.993001
2502 1 360 347 368 360	7 5 5 5 5 5 5 5 5	353 380 357		99.13043 9 104.0346 1 91.25683 8	110.6583 116.0883 109.5592 109.8462	12/6/02 1.9 4.1 3.1 3.8	10 17.9 8.7 13.1	4 4 5 5 5 5 5 4	25/5/02 348.5 3.1 336.5 5.6 345 8.7	12/6/02 3.225 0.48 12.425 2.04 6.175 2.04	26/6/02 103,6636 2,503699 94,96315 1,840356 104,8583 0,993001
1/6/02 1 378 348 364 356	354 337 337	38.7.28		97.3913 9 103.7464 1 92.0765 9	117.03718 1 117.0347 1 109.0141 1	11/6/02			5/02 1.936492 4.784959 6.209871		39869 94745 67427
374 374 348 362 ·	352 342 365 327	3 8 8 8		98.13043 105.1873 105.34426	109.4044 1 116.4036 1 107.6873 1 108 1	10/6/02			2405/02 338.5 1.8 334.25 4.7 341.75 6.2	702 0.686355 1.650676 1.850676	24/05/02 101,2447 1.9 64,33792 1.7 103,9535 1.1
377 377 347 359 359	339 339 319	88.38 88.38		88,26087 8 106,3401 1 87,15847 8	109.0909 1 115.7729 1 109.0141 1 107.3848	9/6/02 3.2 3.2 6.3 5.9 2.9	18.2 17.3 16.1	3.3.3	3.188307 3.520772 7.848177	9/6/02 4.575 0.0 15.4 1.1 4.175 1.1	48256 80375
. 370 370 364 356	325 322 322	385 385		97.97101 8 101.4409 1 87.97614 8	109,4044 1 115,142 1 107,3239 1 107,3846 1	8/6/02			23/05/02 334.25 3.16 331.25 3.52 338.75 7.84		23/05/02 98:9628 1.6 93.4684 0.9 103:0049 0.6
7602 1 367 353 353	353 347 323	351 379 346	7/6/02 103.0899 105.0886 108.6154 109.5678 98.05556	97.10145 9 100 1 88.25137 8	114.8285 106.7606 108.4615	7/6/02			5/02 5.727712 5.93343 6.398614	02 0.800381 1.751844 1.751844	1.387323 1.387323 1.784966 1.286485
75/02 7 384 384 350 350	335 336 338	354 374 348	6/6/02 102.2472 103.2934 106.3333 100.2778	96,23168 9: 103,7464 91,53005 8:	114,6265 1 114,6265 1 105,3521 1 107,0789 1	6/6/02 3.5 7.3 4.9 4.6	17.6 15.7 9.6 16.6	5.5 4.5 4.5	22/06/02 332.25 3.7; 333.75 5.6 334.5 6.3	6/6/02 5.075 0.1 14,925 1.	22/06/02 99.33562 1.3 94.17824 1.7 101.7479 1.2
264 264 360 380 380	352 335 332	£38£3		96.51159 96 102.3055 10 90.71038 91	14,6265 114,6265 106,4769 107,0769	6/6/02		•	5/02 4.25 6.408003 6.060459		54317 45078 14507
385 385 384 384 384	2 2 2 3 3 3 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	8 2 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5		95.65217 96 99.42363 10 94.53552 90	109,4044 10 113,8801 11 107,3239 10 109,5385 10	476/02			2105/02 331.25 4. 331.75 6.40 331.25 6.06	4.252177 4.02707 4.02707	21/05/02 99.03391 1.4: 93.59532 1.5 100.7616 1.2
374 378 388 389	4 2 3 3 5 4 5 4 5 4 5 5 5 5 5 5 5 5 5 5 5 5	¥ 26 8 ¥		97.97101 90 104.611 90 94.80874 94	107.837 10 113.8801 1 104.2254 11 107.3346 11	22.01 5.8 6.4	26.2 16.3 6.6 18.2	7.1 6.4 10.7	.8748 85527 11425	3/6/10 9,4025 16,825 7,625	02 3.643178 1.905917 1.07714
266 366 340 359 351	355 330 348	343 379 348		95.65217 97 104.3228 1 94.53552 94	107.5235 1 113.2492 11 106.7606 16 107.0769 16	2/6/01			20/05/02 320.25 7.4 333 8.11		2005/02 95.86561 3.643178 93.83372 1.905817 100.8764 1.07714
25.02 25.02 25.03	35.25	22.356.25	105.0562 10 105.0562 10 104.1916 10 110.7892 11 107.716 10	63,04346 95 102,5937 10 91,80328 94	107.5235 10 110.4101 11 105.0704 10 105.2308 10	1/6/02			41322 80703 59462	31/05/02 35 0.595119 325 1.348214 725 1.346214	22 237289 263975 863975
375 11 375 3 337 3 354 3 354 3	* # # # # # # # # # # # # # # # # # # #	38 88 88		98.23188 63. 104.611 10; 92.69617 91.	103.3846 10	31/05/02 1 3.4.7 3.9 5.8	119 124 124	2,5 8,0 8,0 8,0	19/05/02 342.5 4.9 338.25 7.11 337.75 8.4	31/05/0 4.35 0. 13.325 1. 6.025 1.	19705/02 102.4294 2.237289 95.4232 1.683975 102.6798 0.659677
30.05/02 31/0 363 3 347 3 355 33	2020	2222		93.62319 96.3 103.17 10 92.62295 92.0	105.0157 108 106.2019 110 103.3603 102 105.5385 103	30/05/02 31/				<b>.</b> -	12 209755 11 252153 9 628252 10
29/05/02 30/0 345 34 345 34 345 34	2222	2222		95.65217 93.6 100 10 94.2623 92.6	103,7616 105 106,8328 108 101,8716 103 105,5365 105	29/05/02 30/		•	16/05/02 327.75 3.275541 329 6.02774 330.333 9.40491	5/02 1,325 1,646461 1,646461	1.311085 97.28794 1.209755 1.311085 97.28794 1.209755 1.406679 92.81281 1.252153 1.000532 99.03276 0.626252
8		2222	28/05/02 29/0 69,7191 101. 102,6946 103. 107,3846 107. 105,2469 106.	93.04346 95.6 101.4409 11 93.1694 B4.2	10,7022 103, 110,0946 108, 102,8169 101, 104,6154 105,	28/05/02 29/0 4.1 9.2 4.0 8.9	18.2 16.9 10.4	84 84 84 84 84	6/02 4.643544 32 8.391464 3 6.342847 330	28/05/02 6.775 1.3 14.85 1.64 9.1 1.64	2 311085 97 199679 82 200532 99
i kreg	% 22 2 X	2222	28.7 89.7 102.1 103.1 105.2 105.2	93.0 101.	\$ 5 5 5	2 4 2 4 8			202 4 4 5 5 5 4 5 4	<b>6</b> ÷ ·	202

5
12/1 363.75 345.25 367
02 6.757712 6.046693 7.553972
11/6/02 361 6.757712 346.75 6.046693 367.25 7.553972
02 5.735489 8.025899 8.041559
10/6/02 360.75 5.7 346.5 8.0 363 8.0
••
USr02 6.480741 10.68 9.18672
359 359 1 345.25 7 362.75
6.25 6.35053 7.659417
358.25 339 361
366502 652087 382412
7/6/02 357 3.306502 339.5 6.652087 360 7.382412
6/6/02 352.5 4.051749 347 7.620812 359.25 8.019067
5.862877 5.86343 7.364048
6/6/ 356.5 343.25 359.25
02 7.011895 4.527693 6.572348
353 343 341.75
4.37975 5.29371 5.87367
363 4.3 363 4.3 350.5 5.2 356 5.5
25.02 5.95119 5.6.90883 9.073772
2/6 3 354.5 9 348.25 3 352
1/6/02 357.75 6.060459 340 7.538789 352 7.22265
357.75 340 352
22 778175 .569132 .238322
31/05/02 355 7,778/75 345.5 6.589/32 347.5 6.238322
502 3.41565 7.157455 6.928203
3.4° 7.15 6.92

2 1/06/02 (2hr) 2 1/06/02 (6hr) 4.215 0.13002 3.076 0.58/1869 12.75 1.57/0233 0.7 2.444381 3.9 1.57/0293 5.85 2.444381 1801 11/04.02 1/00.02


21/06/02 366.25 4.802343 341.5 8.331668 371 7.449632 ND2 1300402 1405402 1505402 1506402 1700402 1505602 1905602 2005602
6.02272 361,75 5.462026 362.5 5.77204 363.25 6.574889 359.25 5.452446 364.75 5.35969 364.5 5.704165 366.75 6.606025 369 6.515202
5.677074 339.5 6.230222 341.25 6.370233 341.25 6.320607 342.5 6.421203 339.75 6.403747 338.75 12,64489 334.25 6.872946
6.296569 366 8.236099 370.5 6.770647 387 7.22285 386.5 7.93203 370.25 6.235079 377 7.322412 368.75 6.774646 374 7.591987

Effect of MD-960 (Smg/kg; s.c. or i.p.) and/or insulin (2<U; s.c.) on body weight (g) of straptozotocin-diabatic rats



- → insulin (s.c.; OU) (n=4)
- -- inculin (s.c.; 2U) (n=4)
- ← LAD-960 ( Smg/lig; s.c. or l.p.) + insulin (2U; s.c.) (n=4)
- ് blud group roselved hould (ഷുറ പു and ലിട്ടോ (Bmgng; arai)
- ජලාල් ලැබුණු අත්තාන් ජලාදිම (Cangle (p.)


Effect of MD-960 (5mg/kg; s.c., i.p. or oral) and/or insulin (244; s.c.) on body weight (% of original) of streptozotocindiabotic rats



- ->- insulin (CU; s.c.) (n=4)
- ->- insulin (2U; s.c.) (n=4)
- whis group roccies insulin MD-230 (Bingis; ora)
- green group received MO-CSD (Emglig; 1.0.)


\*Note: MD-960 represents Gibberellin A3

Effect of MD-960 (Smg/kg; s.c. or I.p.) and/or insulin (2-4U; s.c.) on blood glucose (mM) of streptozotocin-diabetic rats 2 hours after administration



- --- Insulin (4U; s.c.) (n=4)
- -0- Insulin (211; s.c.) (n=4)
- ™D-960 (Smg/kg; s.c. or i.p.) + Insulin (2V; s.c.) (n=4)
- orognoroup rochos CIO-CO (Cincing: 1.0.)
- blue group received CID-CII (Emplig; oral)

Effect of MD-960 (Smg/kg; s.c., i.p. or oral) and/or insulin (2-4U; s.c.) on blood glucose (mM) of streptozotocin-diabetic rats 5 hours after administration



- -- insulin (au; s.c.)
- → MD-960 (5mg/kg; oral) \*/or insulin (2U; s.c.)
- - blue group roceived MDC39 (Emg(kg); crcl)

\*Note: MD-960 represents Gibberellin A<sub>3</sub>

## What is claimed is:

1. A method of treatment comprising adhibiting compounds of formula (1) (Gibberellins) and their pharmaceutically acceptable derivatives for diabetes and related conditions.

#### wherein

 $R^1$  is H or a group  $-O-R^{20}$ , where  $R^{20}$  is H, or together with  $R^2$  or  $R^{10}$  forms a bond  $(C_1-C_2$  or  $C_1-C_{10}$  double bond, respectively);

 $R^2$  is H or a group  $-O-R^{21}$ , where  $R^{21}$  is H, a glycosylic ether group (glycoside ether) or together with  $R^4$  forms a bond (lactone) or together with  $R^1$  or  $R^3$  forms a bond ( $C_1-C_2$  or  $C_2-C_3$  double bond, respectively);

 $R^3$  is H, =O, or -O- $R^{22}$ , where  $R^{22}$  is H or a glycosylic ether group (glycoside ether), or together with  $R^2$  forms a bond ( $C_2$ - $C_3$  double bond);

 $R^4$  is OH, or  $-OR^{23}$ , where  $R^{23}$  is unsubstituted or substituted  $C_{1\sim 20}$  alkyl, allyl, aryl, arylalkyl, amidine,  $-NR^{24}R^{25}$  or an unsaturated or saturated ring containing one or more hetero-atoms selected from the group consisting of nitrogen, oxygen and sulfur;  $R^{24}$  and  $R^{25}$  may or may not be the same, are hydrogen, or  $C_{1\sim 20}$  alkyl, allyl, aryl, arylalkyl or an unsaturated or saturated ring containing one or more hetero-atoms slected from the group consisting of nitrogen, oxygen and sulphur;  $R^4$  together with  $R^{21}$  or  $R^{28}$  forms a bond (lactone);

 $R^5$  is H or a glycosylic ester (glycoside ester) group, or unsubstituted or substituted (e.g. haloginated)  $C_{1\sim20}$  alkyl esters, allyl esters, aryl esters, arylalkyl esters, active esters (such as phenacyl ester, pivaloyl ester);

R<sup>6</sup> is H or OH or together with R<sup>7</sup> forms a bond (C<sub>11</sub>-C<sub>12</sub> double bond);

 $R^7$  is H, =O, or  $-OR^{26}$ , where  $R^{26}$  is H or a glycosylic ether group (glycoside ether) or together with  $R^6$  forms a bond ( $C_{11}$ - $C_{12}$  double bond);

 $R^8$  is H, hydroxyl, mercaptan, or halogen (e.g. F, Cl), amino, azido,  $NR^{24}R^{25}$ , unsubstituted or substituted (e.g. halogenated)  $C_{1\sim20}$  alkyl, allyl, aryl, or arylalkyl, or  $-OR^{27}$ , where  $R^{27}$  is a glycosylic ether group (glycoside ether);

R<sup>9</sup> is H or OH, or together with R<sup>15</sup> forms a bond (C<sub>9</sub>-C<sub>15</sub> bond);

 $R^{10}$  is H, CH<sub>3</sub>, CHO, COOH, or a glycosylic ester (glycoside ester) of said COOH, CH<sub>2</sub>O-R<sup>28</sup> or -OR<sup>28</sup>, where R<sup>28</sup> is H or together with R<sup>4</sup> forms a bond (lactone) or together with R<sup>1</sup> forms a bond (C<sub>1</sub>-C<sub>10</sub> double bond);

R<sup>11</sup> is OH or absent;

R<sup>12</sup> is CH<sub>3</sub>, CH<sub>2</sub>OH, COOH or a glycosylic ester (glycoside ester) of said COOH;

 $R^{13}$  is methylene, or a divalent hetero-atom, or  $NR^{29}$ , where  $R^{29}$  is  $NR^{30}$  or  $OR^{30}$  where  $R^{30}$  is H, or  $C_{1\sim20}$  alkyl, aryl, alkylaryl, when  $R^{11}$  is absent, a double bond is present between  $C_{16}$  and  $R^{13}$ ;

R<sup>14</sup> is H or OH;

R<sup>15</sup> is H, or together with R<sup>9</sup> forms a bond (C<sub>9</sub>-C<sub>15</sub> bond);

And pharmaceutically acceptable derivatives, lactones, esters and salts including alkali metal salts (e.g.  $Na^+$ ,  $K^+$ ), alkaline earth metal salts (e.g.  $Ca^{2+}$ ,  $Mg^{2+}$ ), metal salts (e.g.  $Zn^{2+}$ ,  $Al^{3+}$ ), and salts of ammonium, organic bases (such as  $NR^{16}R^{17}R^{18}R^{19}$  where  $R^{16}$ ,  $R^{17}$ ,  $R^{18}$ ,  $R^{19}$ , which may be the same or not the same, are hydrogen,  $C_{1\sim 20}$  alkyl, alkanol, arryl) thereof.

- 2. The method of claim 1, wherein the Gibberellins are Gibberellin A<sub>3</sub>.
- 3. The method of claim 1, wherein the Gibberellins are a mixture of Gibberellin A<sub>3</sub> and Gibberellin A<sub>4</sub> and/or Gibberellin A<sub>7</sub>.

- 4. The method of claim 1, wherein the pharmaceutically acceptable derivatives are salts including alkali metal salts, alkaline earth metal salts, metal salts and salts of ammonium, organic bases.
- 5. The method of claim 4, wherein the organic bases are NR<sup>16</sup> R<sup>17</sup> R<sup>18</sup> R<sup>19</sup>, where R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup>, R<sup>19</sup>, which may be the same or not the same, are hydrogen, substituted or unsubstituted  $C_{1\sim 20}$  alkyl, alkanol, aryl.
- 6. The method of claim 1, wherein the pharmaceutically acceptable derivatives are lactones, glycoside, esters and active esters.
- 7. A method of treatment of diabetes and related conditions comprising administering an effective amount of a compound of formula (1) (Gibberellins) and their pharmaceutically acceptable derivatives.
- 8. A method according to claim 7, wherein the Gibberellins are Gibberellin A<sub>3</sub>.
- 9. A method according to claim 7, wherein the Gibberellins are a mixture of Gibberellin A<sub>3</sub> and Gibberellin A<sub>4</sub> and/or Gibberellin A<sub>7</sub>.
- 10. A pharmaceutical composition comprising a compound of formula (1), as an active ingredient for the treatment of diabetes and related conditions.
- 11. A slow releasing or long acting pharmaceutical composition comprising a compound of formula (1) as an active ingredient for the treatment of diabetes and related conditions.
- 12. According to claim 11, wherein the formulation is for oral administration.
- 13. According to claim 11, wherein the formulation is for inhalation administration.
- 14. According to claim 11, wherein the formulation is for transdermal administration.
- 15. According to claim 11, wherein the formulation is for injection.
- 16. The method of treatment of diabetes and related conditions comprising compounds of formula (1) (Gibberellins) and their pharmaceutically acceptable derivatives in combination therapy with insulin and/or its fragment derivatives, and/or IGF, and/or growth factors, and/or other pharmaceutically compatible anti-diabetic agents.
- 17. The method according to claim 1, for the treatment of type 1 diabetes.
- 18. The method according to claim 1, for the treatment of type 2 diabetes.

- 19. The method according to claim 1, for the treatment of insulin resistant diabetes.
- 20. The method according to claim 1, for the treatment of diabetic related conditions including obesity, heart and eye diseases, and diabetic ulcers.

**Inventors:** Peter James Jenkins

Minne Wu

David Shine Wu

Australian Biomedical Company Pty. Ltd.

12<sup>th</sup> July, 2002

7 pages of animal Experiment dator is attached.